



Harrison, C. J. (2016). Auxin transport in the evolution of branching forms. *New Phytologist*. <https://doi.org/10.1111/nph.14333>

Peer reviewed version

License (if available):
CC BY-NC

Link to published version (if available):
[10.1111/nph.14333](https://doi.org/10.1111/nph.14333)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Wiley at <http://onlinelibrary.wiley.com/doi/10.1111/nph.14333/full>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Auxin transport in the evolution of branching forms.

C. Jill Harrison[†]

[†]Author for correspondence: jill.harrison@bristol.ac.uk, tel: +44 39411890117

**Present address: School of Biological Sciences. University of Bristol, 24 Tyndall Avenue, Bristol, BS8
1TQ, UK.**

Summary: 117 words

Main text: 2351 words

Number of figures: 1 colour figure

Number of tables: 1

Glossary: 1

13 **Contents list:**

14 Summary

15 I. Diversification of branching forms in land plants

16 II. Roles for auxin transport in branching

17 III. Mechanisms for auxin transport in branching

18 IV. PIN-mediated polar auxin transport is a conserved regulator of branching in vascular plants

19 V. Axial auxin transport is non-polar or weakly polar in liverwort and hornwort sporophytes and bryophyte 20 gametophytes

21 VI. Apical dominance without PIN-mediated polar auxin transport in a moss

22 VII. Conclusions and future perspectives.

23 **Summary**

24 Branching is one of the most striking aspects of land plant architecture, affecting resource acquisition and
25 yield. Polar auxin transport by PIN proteins is a primary determinant of flowering plant branching patterns
26 regulating both branch initiation and branch outgrowth. Several lines of experimental evidence suggest
27 that PIN-mediated polar auxin transport is a conserved regulator of branching in vascular plant
28 sporophytes. However, the mechanisms of branching and auxin transport and relationships between the
29 two are not well known outside the flowering plants and the paradigm for PIN-regulated branching in
30 flowering plants does not fit bryophyte gametophytes. The evidence reviewed here suggests that divergent
31 auxin transport routes contributed to the diversification of branching forms in distinct land plant lineages.

32 **Keywords:** land plant evolution, evo-devo, branching, auxin transport, PIN

33 **I. Diversification of branching forms in land plants**

34 Branching forms have evolved multiple times across the tree of life to optimise exchange between
35 organisms and their environment. In land plants, shoot branching serves two main purposes (1) to optimize
36 photosynthetic efficiency during indeterminate growth (Niklas & Kerchner, 1984) and (2) to determine
37 reproductive output by affecting the number and position of **sporangia** (Niklas, 2004). Branching thus
38 affects plant fitness and yield. Land plants have biphasic life cycles in which a **gametophyte** phase
39 alternates with a **sporophyte** phase, and branching forms have arisen with independent evolutionary
40 trajectories in each life cycle stage (Figure 1) (Harrison, 2016). Current hypotheses of plant interrelationship
41 suggest that vascular plants emerged as a **monophyletic** group from an ancestor shared with bryophytes,
42 but the relationship of bryophytes to vascular plants is contested- bryophytes either form a **paraphyletic**
43 grade at the base of the land plant tree of life (Wickett *et al.*, 2014), or form a monophyletic sister group to
44 vascular plants (Figure 1) (Wickett *et al.*, 2014; Cox *et al.*, 2014). These alternative topologies bear on

45 inferences of the number and direction of branching innovations in land plant evolution in both life cycle
46 stages.

47 A general trend in land plant evolution has been the progressive elaboration of the sporophyte stage of the
48 life cycle at the expense of the gametophyte stage of the life cycle, and sporophytic branching innovations
49 may have contributed to this trend (Niklas, 1997; Niklas, 2004). Whilst the sporophytes of bryophytes are
50 small, comprising a single axis that terminates in sporangium formation, the sporophytes of vascular plants
51 can attain massive sizes as an outcome of branching and indeterminate growth (Figure 1C) (Harrison,
52 2016). Two fossils have forms that are intermediate between living bryophytes and vascular plants.
53 *Partitatheca* is a non-vascular fossil with tiny sporophytes that have stomata and branch (Figure 1E), a
54 character combination suggesting a close affinity to bryophytes (Edwards *et al.*, 2014) and *Cooksonia* fossils
55 reiterate the *Partitatheca* branching pattern but have vascular tissue (Boyce, 2009). These fossils suggest
56 potential steps in the elaboration of sporophyte branching (Figure 1). The number and direction of
57 branching innovations in gametophyte evolution is not yet clear, save to say that there have been multiple
58 origins of branching axial forms in bryophytes and monilophytes, and probably many reversals.

59 II. Roles for auxin transport in branching

60 The mechanisms regulating branching are relatively well characterised in flowering plants, and transport of
61 the plant hormone auxin plays a key role (Domagalska & Leyser, 2011). This was first demonstrated by
62 surgical decapitation experiments in which the shoot apical **meristem** of *Vicia faba* was excised, a
63 manipulation that allows buds initiated in leaf axils to grow out as branches (Thimann & Skoog, 1933). If
64 excision was followed by replacement of the meristem with an agar block impregnated in auxin, the release
65 from suppression did not occur. This experiment suggested that auxin made at the shoot tip can move
66 through a plant to suppress branch outgrowth, a phenomenon known as apical dominance (Thimann &
67 Skoog, 1933). A requirement for long-range **basipetal** polar auxin transport was demonstrated by
68 application of auxin transport inhibitors to a region of the stem, and the subsequent outgrowth of branches
69 below the site of application (Panigrahi & Audus, 1966). More recent work has demonstrated that auxin
70 transport is also required for branch initiation, a process that is intimately linked to leaf initiation (Galweiler
71 *et al.*, 1998; Blakeslee *et al.*, 2007; Bainbridge *et al.*, 2008; Wang, Q *et al.*, 2014; Wang, Y *et al.*, 2014). As
72 leaf **primordia** start to grow out from the shoot apical meristem, an auxin minimum forms in the crease
73 between the new leaf and the shoot apical meristem, and axillary branch meristems form in this region
74 (Wang, Q *et al.*, 2014; Wang, Y *et al.*, 2014).

75

III. Mechanisms for auxin transport in branching

Both the branch initiation and outgrowth processes above are dependent on **PIN** proteins, a class of auxin efflux facilitator with diverse roles in plant development (Petrasek & Friml, 2009). PINs generate directional auxin transport by inserting asymmetrically into plasma membranes (Galweiler *et al.*, 1998). PINs play a key role in leaf initiation by directing auxin to foci on the dome of the shoot apical meristem to determine the position of leaf initiation (Galweiler *et al.*, 1998; Reinhardt *et al.*, 2000). As leaf primordia start to grow out PINs direct auxin transport away from the crease forming between the shoot apical meristem and the new leaf, leading to the formation of auxin minima and axillary meristem initiation (Wang, Q *et al.*, 2014; Wang, Y *et al.*, 2014). PINs play a key role in branch outgrowth by generating and modulating long-range basipetal auxin transport in the stem as well as modulating the action of other hormonal cues on branching (Okada *et al.*, 1991; Shinohara *et al.*, 2013; Bennett *et al.*, 2016). **ABCB/PGP** auxin efflux transporters also effect long-range basipetal transport and modulate branch outgrowth, acting synergistically with PINs (Noh *et al.*, 2001; Blakeslee *et al.*, 2007), and **AUX/LAX** influx symporters effect auxin influx in leaf initiation, thereby modulating branching (Swarup *et al.*, 2004; Yang *et al.*, 2006; Bainbridge *et al.*, 2008). Cell to cell connectivity via **plasmodesmata** has recently been identified as a further potential route for auxin transport in the regulation of plant development (Han *et al.*, 2014), and deposition or break down of the cell wall polymer callose by **CalS** and **GHL** genes is one way that the size of plasmodesmatal openings can be actively regulated. All aforementioned families are conserved within the land plants and are therefore potential contributors to auxin transport in the evolution of branching forms (Del Bem & Vincentz, 2010; Carraro *et al.*, 2012; Bennett, T *et al.*, 2014; Gaudioso-Pedraza & Benitez-Alfonso, 2014; Lane *et al.*, 2016).

IV. PIN-mediated polar auxin transport is a conserved regulator of branching in vascular plants

The bulk basipetal pattern of auxin transport that regulates **axillary branching** in flowering plants can be measured using radiolabelled auxin transport assays (Figure 1D, Table 1) (Goldsmith, 1966; Goldsmith, 1977). Tritiated auxin is applied to one cut surface of a stem segment, and its passage through the stem is monitored by extracting auxin from a portion at the other end of the segment in scintillation fluid. The rate of auxin transport is inferred by measuring radioactive decay using a scintillation counter in relation to time allowed for transport (Lewis & Muday, 2009). Such assays demonstrate that bulk basipetal auxin transport is a conserved property of vascular plant sporophytes (Steeves & Briggs, 1960; Wochok & Sussex, 1973; Wochok & Sussex, 1974; Walters & Osborne, 1979), and **PAT1** sensitivity suggests that PINs provide the transport (Sanders & Langdale, 2013). The branching mechanism of basal vascular plant lineages involves **bifurcation** of the main shoot apex (in lycophytes and some monilophytes) or the establishment of new apices in leaf axils (in monilophytes; Figure 1) (Bierhorst, 1977; Harrison *et al.*, 2007; Schneider, 2012), but links between PIN-mediated **PAT** and branching are yet to be identified. However, decapitation

110 experiments in both lycophytes and monilophytes suggest that there is likely conservation. In the lycophyte
111 *Selaginella*, apical decapitation can result in the outgrowth of angle meristems initiated at branch
112 divergence points, mirroring the PIN-mediated plasticity of branching in flowering plants, and auxin applied
113 at the tip counteracts this effect (Williams, 1937; Seidl, 1941; Jernstedt *et al.*, 1994). Similarly, decapitation
114 can result in bud formation from the rhizome in ferns (Wardlaw, 1965). PATI sensitive basipetal auxin
115 transport can be detected in the single-stemmed axes of moss sporophytes (Poli *et al.*, 2003; Fujita *et al.*,
116 2008), and disruption of *Physcomitrella patens* (*Physcomitrella*) PIN function can induce branching to
117 generate a similar form to the most ancient branching fossils (Fujita *et al.*, 2008; Bennett, TA *et al.*, 2014).
118 Current evidence suggests that PIN-mediated PAT is a conserved regulator of sporophyte branching in land
119 plants.

120 **V. Axial auxin transport is non-polar or weakly polar in liverwort and hornwort** 121 **sporophytes and bryophyte gametophytes**

122 Although moss sporophytes show PATI sensitive bulk basipetal auxin transport (Poli *et al.*, 2003; Fujita *et*
123 *al.*, 2008), other bryophyte sporophytes do not (Figure 1C, D). Transport assays in the hornwort *Phaeoceros*
124 *pearsonii* detect low rates of PATI-insensitive transport in both directions in sporophyte axes, consistent
125 with a diffusive mechanism (Poli *et al.*, 2003). Assays similarly detect low rates of bidirectional transport in
126 the liverwort *Pellia epiphylla*, but transport is PATI-sensitive suggesting the involvement of membrane
127 transporters (Thomas, 1980; Poli *et al.*, 2003). Transport in bryophyte gametophytes can be polar
128 (Maravolo, 1976; Rose & Bopp, 1983; Rose *et al.*, 1983), bidirectional (Fujita *et al.*, 2008; Coudert *et al.*,
129 2015; ^aPiatowski *et al.*, 2012), or have a weak directional bias (Gaal *et al.*, 1982), and transport may or may
130 not be sensitive to PATI and **NOA** inhibitors (Maravolo, 1976; Rose & Bopp, 1983; Fujita *et al.*, 2008;
131 Bennett, TA *et al.*, 2014; ^aPiatowski *et al.*, 2012) that interfere with membrane transporter function
132 (Geldner *et al.*, 2001; Parry *et al.*, 2001). The roles of PIN proteins and other auxin transporters have not yet
133 been characterised in liverworts and hornworts, so the mechanisms for transport in these groups are not
134 clear. In *Physcomitrella* **gametophores** PINs can target to the plasma membrane and are likely to generate
135 short-range directional transport in a manner similar to the pattern observed in flowering plant shoot
136 apices (Bennett, TA *et al.*, 2014). Disruption of PIN function and PATI application induces defects in
137 gametophore apex function and leaf development (Bennett, TA *et al.*, 2014; Viaene *et al.*, 2014), but not
138 gametophore branching (Coudert *et al.*, 2015). In combination, these data suggest that the roles of PIN-
139 mediated auxin transport in branching are not conserved between bryophytes and vascular plants; the
140 moss sporophyte exception discussed above may reflect convergence.

141 **VI. Apical dominance without PIN-mediated polar auxin transport in a moss**

142 Recent work in *Physcomitrella* gametophores suggests an alternative route for auxin transport in the
143 regulation of branching (Coudert *et al.*, 2015). Branches initiate by respecification of epidermal cells into

apical cells in leaf axils after the leaf has developed, and subsequent outgrowth is not delayed (Coudert *et al.*, 2015). Surgical decapitation experiments similar to the experiments undertaken in flowering plant sporophytes showed that the main gametophore apex is an auxin source that acts at a distance to suppress branch initiation (von Maltzahn, 1959; Nyman & Cutter, 1981; Coudert *et al.*, 2015). *Physcomitrella pinab* mutants initiate branches in a fairly normal pattern, and although there are two other PINs in *Physcomitrella*, application of **NPA** to *pinab* mutants does not disrupt branching further (Coudert *et al.*, 2015). These results suggest that PINs do not provide the auxin transport required to regulate branch initiation, and a computational modelling approach showed that the measured bi-directional property of auxin transport in moss gametophores is required to attain realistic branch initiation patterns (Coudert *et al.*, 2015). Modelling also predicted that variation in the rate of bidirectional auxin transport should perturb branch initiation patterns, and pharmacological treatments with the callose synthesis inhibitor **DDG** (Jaffe & Leopold, 1984) perturbed initiation patterns in a similar way to model predictions (Coudert *et al.*, 2015). These observations point to a role for auxin transport via plasmodesmata in branching in a moss, but links between auxin transport and branching have not yet been interrogated in other bryophytes.

VII. Conclusions and future perspectives

As yet there is very scant data relating to the auxin transport properties of non flowering plants. The bryophyte data discussed here were obtained by different labs and have used different taxon sampling and techniques. In some instances experiments with the same species or different species within the same lineage have generated different results (Maravolo, 1976; Gaal *et al.*, 1982; Poli *et al.*, 2003; ^aPiatowski *et al.*, 2012). Branching patterns and their developmental basis are also poorly characterised in non-flowering plants (most notably in monilophytes), and land plant phylogeny remains contested at key nodes for inferences relating to branching evolution (Cox *et al.*, 2014; Wickett *et al.*, 2014). Despite these deficiencies in our knowledge, the data discussed suggest that divergent auxin transport pathways were recruited to regulate branching in land plant gametophytes and sporophytes during evolution. Current evidence suggests that PIN-mediated basipetal auxin transport is a conserved mechanism within vascular plants (Sanders & Langdale, 2013), and a conserved regulator of branching. Recent work shows that there is also bulk basipetal auxin transport in moss sporophytes, and demonstrates that perturbing PIN function can induce bifurcation (Fujita *et al.*, 2008; Bennett, TA *et al.*, 2014). In contrast bi-directional auxin transport regulates gametophytic branching patterns, potentially via plasmodesmata (Coudert *et al.*, 2015). The nature of auxin transporters in liverworts and hornworts, the ancestral mechanism for auxin transport within land plants, and links between auxin transport and branching in bryophyte gametophytes are open questions that are now amenable to experimental manipulation.

Acknowledgements

177 I would like to thank Jane Langdale, Harald Schneider and two anonymous referees for helpful comments
178 on manuscript drafts, and staff at the New Phytologist for help with manuscript preparation. I am grateful
179 to the Royal Society for funding my work.

References

- Albaum HG. 1938. Inhibitions due to growth hormones in fern prothallia and sporophytes. *American Journal of Botany* 25: 124-133.
- Bainbridge K, Guyomarc'h S, Bayer E, Swarup R, Bennett M, Mandel T, Kuhlemeier C. 2008. Auxin influx carriers stabilize phyllotactic patterning. *Genes & development* 22(6): 810-823.
- Bennett MJ, Marchant A, Green HG, May ST, Ward SP, Millner PA, Walker AR, Schulz B, Feldmann KA. 1996. *Arabidopsis AUX1* gene: a permease-like regulator of root gravitropism. *Science* 273: 948-950.
- Bennett T, Brockington S, Rothfels C, Graham S, Stevenson D, Kutchan T, Rolf M, Thomas P, Wong GK-S, Leyser O, Glover BJ, Harrison CJ. 2014. Paralagous radiations of PIN proteins with multiple origins of non-canonical PIN structure. *Molecular Biology and Evolution* 31(8): 2042-2060.
- Bennett T, Hines G, van Rongen M, Waldie T, Sawchuk MG, Scarpella E, Ljung K, Leyser O. 2016. Connective auxin transport in the shoot facilitates communication between shoot apices. *PLoS Biol* 14(4): e1002446.
- Bennett TA, Liu MM, Aoyama T, Bierfreund NM, Braun M, Coudert Y, Dennis RJ, O'Connor D, Wang XY, White CD, Decker EL, Reski R, Harrison CJ. 2014. Plasma membrane-targeted PIN proteins drive shoot development in a moss. *Current Biology* 24: 2776-2785.
- Berthier J. 1973. Recherches sur la structure et le développement de l'apex du gamétophyte feuillé des mousses. *Revue bryologique et lichenologique* 38: 421-551.
- Bierhorst DW. 1977. On the stem apex, leaf initiation and early leaf ontogeny in filiclean ferns. *American Journal of Botany* 64: 125-152.
- Blakeslee JJ, Bandyopadhyay A, Lee OR, Mravec J, Titapiwatanakun B, Sauer M, Makam SN, Cheng Y, Bouchard R, Adamec J, Geisler M, Nagashima A, Sakai T, Martinoia E, Friml J, Peer WA, Murphy AS. 2007. Interactions among PIN-FORMED and P-glycoprotein auxin transporters in *Arabidopsis*. *The Plant Cell* 19(1): 131-147.
- Boyce C. 2009. How green was *Cooksonia*? The importance of size in understanding the early evolution of physiology in the vascular plant lineage. *Paleobiology* 34: 179-194.
- Buchloh G. 1951. *Symmetrie und verzweigung der lebermoose*. Heidelberg: Springer-Verlag.
- Carraro N, Tisdale-Orr TE, Clouse RM, Knöller AS, Spicer R. 2012. Diversification and expression of the *PIN*, *AUX/LAX*, and *ABCB* families of putative auxin transporters in *Populus*. *Frontiers in Plant Science* 3: 17.
- Cho M, Lee SH, Cho HT. 2007. P-glycoprotein4 displays auxin efflux transporter-like action in *Arabidopsis* root hair cells and tobacco cells. *Plant Cell* 19: 3930-3943.
- Coudert Y, Palubicki W, Ljung K, Novak O, Leyser O, Harrison CJ. 2015. Three ancient hormonal cues coordinate shoot branching in a moss. *eLIFE* 4: e06808.

214 Cox CJ, Li B, Foster PG, Embley M, Civiš P. 2014. Conflicting phylogenies for early land plants are caused by
215 composition biases among synonymous substitutions. *Systematic Biology* 63: 272-279.

216 Crandall-Stotler B. 1972. Morphogenetic patterns of branch formation in the leafy hepaticae- a resumé. *The*
217 *Bryologist* 75: 381-403.

218 Del Bem LE, Vincentz MG. 2010. Evolution of xyloglucan-related genes in green plants. *BMC Evolutionary*
219 *Biology* 10(1): 341.

220 Ding Z, Wang B, Moreno I, Dupláková N, Simon S, Carraro N, Reemmer J, Pěnčík A, Chen X, Tejos R, Skůpa P,
221 Pollmann S, Mravec J, Petrášek J, Zažímalová E, Honys D, Rolčík J, Murphy A, Orellana A, Geisler M,
222 Friml J. 2012. ER-localized auxin transporter PIN8 regulates auxin homeostasis and male
223 gametophyte development in *Arabidopsis*. *Nature Communications* 3: 941.

224 Domagalska MA, Leyser O. 2011. Signal integration in the control of shoot branching. *Nat Rev Mol Cell Biol*
225 12: 211-221.

226 Edwards D, Morris JL, Richardson JB, Kenrick P. 2014. Cryptospores and cryptophytes reveal hidden
227 diversity in early land floras. *New Phytologist* 202: 50-78.

228 Fujita T, Sakaguchi H, Hiwatashi Y, Wagstaff SJ, Ito M, Deguchi H, Sato T, Hasebe M. 2008. Convergent
229 evolution of shoots in land plants: lack of auxin polar transport in moss shoots. *Evolution and*
230 *Development* 10: 176-186.

231 Gaal DJ, Dufresne SJ, Maravolo NC. 1982. Transport of ¹⁴C-indoleacetic acid in the hepatic *Marchantia*
232 *polymorpha*. *The Bryologist* 85: 410-418.

233 Galweiler L, Guan C, Muller A, Wisman E, Mendgen K, Yephremov A, Palme K. 1998. Regulation of polar
234 auxin transport by AtPIN1 in *Arabidopsis* vascular tissue. *Science* 282: 2226-2230.

235 Gaudioso-Pedraza R, Benitez-Alfonso Y. 2014. A phylogenetic approach to study the origin and evolution of
236 plasmodesmata-localized *glycosyl hydrolases family 17*. *Frontiers in Plant Science* 5: 212.

237 Geisler M, Blakeslee JJ, Bouchard R, Lee OR, Vincenzetti V, Bandyopadhyay A, Titapiwatanakun B, Peer WA,
238 Bailly A, Richards EL, Ejendal KF, Smith AP, Baroux C, Grossniklaus U, Muller A, Hrycyna CA, Dudler
239 R, Murphy AS, Martinoia E. 2005. Cellular efflux of auxin catalyzed by the *Arabidopsis* MDR/PGP
240 transporter AtPGP1. *Plant Journal* 44: 179–194.

241 Geldner N, Friml J, Stierhof YD, Jürgens G, Palme K. 2001. Auxin transport inhibitors block PIN1 cycling and
242 vesicle trafficking. *Nature* 413: 425-428

243 Goldsmith MHM. 1966. Maintenance of polarity of auxin movement by basipetal transport. *Plant*
244 *Physiology* 41: 749-754.

245 Goldsmith MHM. 1977. The polar transport of auxin. *Annual Review of Plant Physiology* 28: 439-478.

246 Gregorich M, Fisher R. 2006. Auxin regulates lateral meristem activation in developing gametophytes of
247 *Ceratopteris richardii*. *Canadian Journal of Botany* 84: 1520-1530.

248 Han X, Hyun TK, Zhang M, Kumar R, Koh EJ, Kang BH, Lucas WJ, Kim JY. 2014. Auxin-callose-mediated
 249 plasmodesmal gating is essential for tropic auxin gradient formation and signaling. *Developmental*
 250 *Cell* 28: 132-146.

251 Harrison CJ. 2015. Shooting through time: new insights from transcriptomic data. *Trends in Plant Science*
 252 20: 468-470.

253 Harrison CJ. 2016. Development and genetics in the evolution of land plant body plans. *Philosophical*
 254 *Transactions of the Royal Society B* In Press.

255 Harrison CJ, Rezvani M, Langdale JA. 2007. Growth from two transient apical initials in the meristem of
 256 *Selaginella kraussiana*. *Development* 134: 881-889.

257 Hou G, Hill JP, Blancaflor EB. 2004. Developmental anatomy and auxin response of lateral root formation in
 258 *Ceratopteris richardii*. *Journal of Experimental Botany* 55: 685-693.

259 Jaffe M, Leopold A. 1984. Callose deposition during gravitropism of *Zea mays* and *Pisum sativum* and its
 260 inhibition by 2-deoxy-D-glucose. *Planta* 161(1): 20-26.

261 Jernstedt JA, Cutter EG, Lu P. 1994. Independence of organogenesis and cell pattern in developing angle
 262 shoots of *Selaginella martensii*. *Annals of Botany* 74: 343-355.

263 Kim JY, Henrichs S, Bailly A, Vincenzetti V, Sovero V, Mancuso S, Pollmann S, Kim D, Geisler M, Nam H-G.
 264 2010. Identification of an ABCB/P-glycoprotein-specific inhibitor of auxin transport by chemical
 265 genomics. *Journal of Biological Chemistry* 285: 23309-23317.

266 La Farge-England C. 1996. Growth form, branching pattern, and perichaetial position in mosses: cladocarp
 267 and pleurocarpy redefined. *The Bryologist* 99: 170-186.

268 Lane TS, Rempe CS, Davitt J, Staton ME, Peng Y, Soltis DE, Melkonian M, Deyholos M, Leebens-Mack JH,
 269 Chase M. 2016. Diversity of ABC transporter genes across the plant kingdom and their potential
 270 utility in biotechnology. *BMC biotechnology* 16(1): 1.

271 Lewis DR, Muday GK. 2009. Measurement of auxin transport in *Arabidopsis thaliana*. *Nature Protocols* 4:
 272 437-451.

273 Maravolo NC. 1976. Polarity and localization of auxin movement in the hepatic, *Marchantia polymorpha*.
 274 *American Journal of Botany* 63: 526-531.

275 Nayar BK, Kaur S. 1971. Gametophytes of homosporous ferns. *Botanical Review* 3: 295-396.

276 Niklas KJ. 1997. *The evolutionary biology of plants*. Chicago: University of Chicago Press. P161.

277 Niklas KJ. 2004. Computer models of early land plant evolution. *Annual Review of Earth and Planetary*
 278 *Science*: 32: 47-66.

279 Niklas KJ, Kerchner V. 1984. Mechanical and photosynthetic constraints on the evolution of plant shape.
 280 *Paleobiology* 10: 79-101.

281 Noh B, Murphy AS, Spalding EP. 2001. Multidrug resistance-like genes of *Arabidopsis* required for auxin
 282 transport and auxin-mediated development. *The Plant Cell* 13(11): 2441-2454.

283 Nyman LP, Cutter EG. 1981. Auxin–cytokinin interaction in the inhibition, release, and morphology of
 284 gametophore buds of *Plagiomnium cuspidatum* from apical dominance. *Canadian Journal of Botany*
 285 59: 750-762.

286 Okada K, Ueda J, Komaki MK, Bell CJ, Shimura Y. 1991. Requirement of the auxin polar transport system in
 287 early stages of *Arabidopsis* floral bud formation. *The Plant Cell* 3: 677-684.

288 Panigrahi B, Audus L. 1966. Apical dominance in *Vicia faba*. *Annals of Botany* 30(3): 457-473.

289 Parihar NS. 1967. *Bryophyta*. Allahabad: Indian Universities Press.

290 Parry G, Delbarre A, Marchant A, Swarup R, Napier R, Perrot-Rechenmann C, Bennett MJ. 2001. Novel auxin
 291 transport inhibitors phenocopy the auxin influx carrier mutation *aux1*. *Plant Journal* 25: 399-406.

292 Petrasek J, Friml J. 2009. Auxin transport routes in plant development. *Development* 136: 2675-2688.

293 Piatkowski B, Bader G, Wallace A, Kellam L, Branning J, Corbicz M, Poli D 2012. Examining polar auxin
 294 transport evolution in bryophyte gametophytes, *Botany 2012*. Columbus, Ohio. URL
 295 <http://2012.botanyconference.org/engine/search/index.php?func=detail&aid=502> [accessed 29
 296 September 2016].

297 Poli D, Jacobs M, Cooke TJ. 2003. Auxin regulation of axial growth in bryophyte sporophytes: its potential
 298 significance for the evolution of early land plants. *American Journal of Botany* 90: 1405-1415.

299 Reinhardt D, Mandel T, Kuhlemeier C. 2000. Auxin regulates the initiation and radial position of plant lateral
 300 organs. *The Plant Cell* 12: 507-518.

301 Rose S, Bopp M. 1983. Uptake and polar transport of indoleacetic acid in moss rhizoids. *Physiologica*
 302 *Plantarum* 58: 57-61.

303 Rose S, Rubery PH, Bopp M. 1983. The mechanism of auxin uptake and accumulation in moss protonemata.
 304 *Physiologia Plantarum* 58(1): 52-56.

305 Sanders HL, Langdale JA. 2013. Conserved transport mechanisms but distinct auxin responses govern shoot
 306 patterning in *Selaginella kraussiana*. *New Phytologist* 198: 419-428.

307 Schneider H 2012. Evolutionary morphology of ferns (monilophytes). In: Ambrose BA, Purugganan M eds.
 308 *The evolution of plant form*: Wiley-Blackwell.

309 Seidl W. 1941. Wuchsstoffuntersuchungen an *Selaginella martensii*. *Jahrbücher für Wissenschaftlichen*
 310 *Botanik* 89: 832-861.

311 Shinohara N, Taylor C, Leyser O. 2013. Strigolactone can promote or inhibit shoot branching by triggering
 312 rapid depletion of the auxin efflux protein PIN1 from the plasma membrane. *PLOS Biology* 11(1):
 313 e1001474.

314 Steeves TA, Briggs WR. 1960. Morphogenetic studies on *Osmunda cinnamomea* L. The auxin relationships
 315 of expanding fronds. *Journal of Experimental Botany* 11: 45-67.

316 Swarup R, Kargul J, Marchant A, Zadik D, Rahman A, Mills R, Yemm A, May S, Williams L, Millner P. 2004.
 317 Structure-function analysis of the presumptive *Arabidopsis* auxin permease AUX1. *The Plant Cell*
 318 16(11): 3069-3083.

319 Thimann KV, Skoog F. 1933. Studies on the growth hormone of plants: III. The inhibiting action of the
 320 growth substance on bud development. *PNAS* 19: 714-716.
 321 Thomas RJ. 1980. Cell elongation in hepatics: the seta system. *Bulletin of the Torrey Botanical Club* 107:
 322 339-345.
 323 Viaene T, Landberg K, Thelander M, Medvecka E, Pederson E, Feraru E, Cooper ED, Karimi M, Delwiche CF,
 324 Ljung K, Geisler M, Sundberg E, Friml J. 2014. Directional auxin transport mechanisms in early
 325 diverging land plants. *Current Biology* 24: 2786-2791.
 326 von Maltzahn KE. 1959. Interaction between kinetin and indoleacetic acid in the control of bud reactivation
 327 in *Splachnum ampullaceum* (L.) Hedw. *Nature* 183: 60-61.
 328 Walters J, Osborne DJ. 1979. Ethylene and auxin-induced cell growth in relation to auxin transport and
 329 metabolism and ethylene production in the semi-aquatic plant, *Regnellidium diphyllum*. *Planta* 146:
 330 309-317.
 331 Wang Q, Kohlen W, Rossmann S, Vernoux T, Theres K. 2014. Auxin depletion from the leaf axil conditions
 332 competence for axillary meristem formation in *Arabidopsis* and tomato. *The Plant Cell* 26: 2068-
 333 2079.
 334 Wang Y, Wang J, Shi B, Yu T, Qi J, Meyerowitz EM, Jiao Y. 2014. The stem cell niche in leaf axils is
 335 established by auxin and cytokinin in *Arabidopsis*. *The Plant Cell* 26: 2055-2067.
 336 Wardlaw CW 1965. The organization of the shoot apex. In: Lang A ed. *Differenzierung und*
 337 *Entwicklung/Differentiation and Development*: Springer-Verlag Berlin Heidelberg. 966-1076.
 338 Wickett NJ, Mirarab S, Nguyen N, Warnow T, Carpenter E, Matascie N, Ayyampalayam S, Barker MS,
 339 Burleigh JG, Gitzendanner MA, Ruhfel BR, Wafula E, Der JP, Graham SW, Mathews S, Melkonian M,
 340 Soltis DE, Soltis PS, Miles NW, Rothfels CJ, Pokorný L, Shaw AJ, DeGironimo L, Stevenson DW, Surek
 341 B, Villarreal JC, Roure B, Philippe H, de Pamphilis CW, Chen T, Deyholos MK, Baucom RS, Kutchan
 342 TM, Augustin MM, Wang J, Zhang Y, Tian Z, Yan Z, Wu X, Sun X, K-S. WG, Leebens-Mack J. 2014.
 343 Phylotranscriptomic analysis of the origin and early diversification of land plants. *PNAS* 111: E4859–
 344 E4868.
 345 Williams S. 1937. Correlation phenomena and hormones in *Selaginella*. *Nature* 139: 966.
 346 Wochok ZS, Sussex IM. 1973. Morphogenesis in *Selaginella* I: auxin transport in the stem. *Plant Physiology*
 347 51: 646-650.
 348 Wochok ZS, Sussex IM. 1974. Morphogenesis in *Selaginella* II: Auxin transport in the root (rhizophore).
 349 *Plant Physiology* 53: 738-741.
 350 Yang Y, Hammes UZ, Taylor CG, Schachtman DP, Nielsen E. 2006. High-affinity auxin transport by the AUX1
 351 influx carrier protein. *Current Biology* 16(11): 1123-1127.
 352
 353
 354

355

356

357

Figure legend

Figure 1: Auxin transport pathways in the evolution of branching forms.

(A) Summary of axial gametophyte forms in bryophytes and monilophytes. Whilst hornwort and liverwort thalli branch dichotomously (Parihar, 1967), leafy liverwort and moss gametophores branch laterally (Buchloh, 1951; von Maltzahn, 1959; Crandall-Stotler, 1972; Berthier, 1973; La Farge-England, 1996; Coudert *et al.*, 2015). Monilophyte prothalli bifurcate or branch laterally (Nayar & Kaur, 1971; Schneider, 2012).

(B) Summary of documented auxin transport properties in bryophyte and monilophyte gametophytes. Whilst hornwort thalli have no detectable basipetal transport (^aPiatowski *et al.*, 2012), liverwort thalli may (Maravolo, 1976) or may not (Gaal *et al.*, 1982; ^aPiatowski *et al.*, 2012) have basipetal transport and transport may or may not be PATI sensitive (Maravolo, 1976; Gaal *et al.*, 1982; ^aPiatowski *et al.*, 2012) and NOA sensitive (^aPiatowski *et al.*, 2012). In moss gametophores the patterns of transport are consistent with bi-directional transport by membrane transporters or plasmodesmata (Fujita *et al.*, 2008; Coudert *et al.*, 2015), and a functional link between auxin transport and branching has been demonstrated in *Physcomitrella* (Coudert *et al.*, 2015). Although there is PIN-mediated auxin transport in *Physcomitrella* gametophores (Bennett, T *et al.*, 2014; Bennett, TA *et al.*, 2014; Viaene *et al.*, 2014), it is a minor contributor to the regulation of branching patterns. Experiments with DDG callose synthesis inhibitors (Jaffe & Leopold, 1984) support the hypothesis that a plasmodesmatal mechanism regulates branching (Coudert *et al.*, 2015). Monilophyte gametophyte development is sensitive to NPA (Gregorich and Fisher, 2006).

(C) Summary of sporophyte forms in land plants. Whilst bryophytes have uni-axial sporophytes that terminate with sporangia (Parihar, 1967), vascular plants branch by bifurcation (Bierhorst, 1977; Harrison *et al.*, 2007) or branch laterally (Domagalska & Leyser, 2011).


(D) Summary of documented sporophytic auxin transport properties in land plants. Hornwort sporophyte axes do not directionally transport auxin and are PATI insensitive suggesting a diffusive mechanism (Poli *et al.*, 2003). Liverwort sporophyte axes similarly lack directionally biased auxin transport, but transport is PATI sensitive (Poli *et al.*, 2003), consistent with a contribution for membrane transporters and/or plasmodesmatal connectivity. Moss sporophytes have bulk basipetal auxin transport (Poli *et al.*, 2003; Fujita *et al.*, 2008) that is PATI (Poli *et al.*, 2003; Fujita *et al.*, 2008) and NOA (Poli *et al.*, 2003) sensitive. Disruption of PIN function causes sporophytic abnormalities including bifurcation (Fujita *et al.*, 2008; Bennett, TA *et al.*, 2014). The data suggest that bulk basipetal transport in mosses is PIN-mediated, and that the innovation of sporophytic branching forms may have involved changes in PIN function (Harrison, 2015). Lycophyte (Wochok & Sussex, 1973; Wochok & Sussex, 1974; Sanders & Langdale, 2013) and monilophyte (Walters & Osborne, 1979) sporophytes have bulk-basipetal transport that is PATI sensitive

392 (Hou *et al.*, 2004; Sanders & Langdale, 2013). In a lycophyte, PATI application disrupts bifurcation (Sanders
393 & Langdale, 2013), but as yet there are no functional data on the nature of auxin transporter involved. In
394 angiosperms, NPA sensitive (Geldner *et al.*, 2001) PIN-mediated bulk basipetal auxin transport is a major
395 determinant of branching patterns (Domagalska & Leyser, 2011) and NOA sensitive AUX/LAX auxin influx
396 carriers (Bennett *et al.*, 1996; Parry *et al.*, 2001; Bainbridge *et al.*, 2008), **BUM** sensitive ABCB/PGP efflux
397 carriers (Noh *et al.*, 2001; Geisler *et al.*, 2005; Blakeslee *et al.*, 2007; Cho *et al.*, 2007; Kim *et al.*, 2010) also
398 regulate branching.

399 (E) Current hypothesis of phylogenetic relationships between major land plant groups (Cox *et al.*, 2014;
400 Wickett *et al.*, 2014) showing potential steps involved in the innovation of branching forms as suggested by
401 the fossil record (Edwards *et al.*, 2014).

402

403 **Figure 1**



404

405

406 **Table 1**

407

408

	Generation	Hornworts	Liverworts	Mosses	Lycophytes	Monilophytes	Angiosperms
1. Bulk basipetal transport	G	- ^a	+/- ^{2,3,a}	- ^{5,a}	n.a.		n.a
	S	- ¹	- ¹	+ ^{1,5}	+/- ¹⁰⁻¹²	+ ¹³	+ ^{1,5,18,19}
2. Directional transport	G		+/- ^{2,3}	+/- ⁵⁻⁹		+ ^{14,15}	
	S	- ¹	- ^{1,4}	+ ^{1,5}	+ ¹⁰⁻¹²	+ ^{13,15,16}	+ ^{1,5,18,19}
3. Polar ratio in axis	G						
	S	1 ¹	1.1 ¹	9.3 ¹	c. 2 ¹⁰	c. 100 ¹³	674 ¹
4. Rate of transport mmh ⁻¹	G		14 ³				
	S		6.9 ¹	8.9 ¹	12 ¹⁰		11 ¹
5. NPA or TIBA sensitivity	G		+ ²	+ ^{7-9,a}		+ ¹⁴	
	S	- ¹	+ ¹	+ ^{1,5}	+ ¹²	+ ¹⁷	+ ^{1,20}
6. PIN-mediated Transport	G			+ ^{8,9}			+ ²¹
	S			+ ⁸			+ ²²
7. NOA sensitivity	G			+ ^a			
	S			+ ¹			+ ^{23,24}

409

410 **Table 1: Summary of auxin transport data from major land plant clades and data sources.**

411 G: gametophyte, S: sporophyte, +: presence, -: absence, blank: no data. Sources: ¹Poli *et al.*, 2003,
412 ²Maravolo, 1976, ³Gaal *et al.*, 1982, ⁴Thomas, 1980, ⁵Fujita *et al.*, 2008, ⁶Rose *et al.*, 1983, ⁷Rose & Bopp,
413 1983, ⁸Bennett, TA *et al.*, 2014, ⁹Viaene *et al.*, 2014, ¹⁰Wochok & Sussex, 1973, ¹¹Wochok & Sussex, 1974,
414 ¹²Sanders & Langdale, 2013, ¹³Walters & Osborne, 1979, ¹⁴Gregorich & Fisher, 2006, ¹⁵Albaum, 1938,
415 ¹⁶Steeves & Briggs, 1960, ¹⁷Hou *et al.*, 2004, ¹⁸Goldsmith, 1966, ¹⁹Goldsmith, 1977, ²⁰Geldner *et al.*, 2001,
416 ²¹Ding *et al.*, 2012, ²²Galweiler *et al.*, 1998, ²³Bennett *et al.*, 1996, ²⁴Parry *et al.*, 2001 ^aPiatowski *et al.*, 2012
417 (online conference abstract).

418

419

420 **Box 1: Glossary**

- 421 **ABCB/PGP:** A class of membrane targeted auxin efflux transporter.
- 422 **AUX/LAX:** A membrane targeted auxin influx symporter.
- 423 **Axillary branching:** Branching that arises due to the activity of meristems in leaf axils.
- 424 **Basipetal:** From the apex to the base.
- 425 **Bifurcation:** Branching that occurs by apex divergence.
- 426 **BUM:** 2-[4-(diethylamino)-2-hydroxybenzoyl]benzoic acid, a compound that inhibits ABCB/PGP-mediated
427 auxin transport
- 428 **CalS/ GSL:** Callose Synthase/ Glucan Synthase enzymes that polymerize glucose to form Callose (β -1,3-
429 Glucan).
- 430 **DDG:** 2-deoxy-D-glucose, a chemical inhibitor of callose synthesis.
- 431 **Gametophore:** A gamete producing leafy shoot in mosses and liverworts.
- 432 **Gametophyte:** A stage of the land plant life cycle during which gamete production occurs.
- 433 **GHL:** Glycosyl hydrolase enzymes that hydrolyze 1,3-beta-glucan polysaccharides such as callose.
- 434 **Meristem:** The growing tip of a plant.
- 435 **Monophyletic group:** A group containing all the species (or genes) derived from a shared ancestral lineage.
- 436 **NOA:** Naphthoxyacetic acid, a compound that inhibits AUX1-mediated auxin influx.
- 437 **NPA:** 1-N-Naphthylphthalamic acid, a compound that inhibits PIN-mediated auxin efflux.
- 438 **Paraphyletic group:** A group of species (or genes) that is not monophyletic and is basal with respect to a
439 single monophyletic group.
- 440 **PAT:** Polar auxin transport.
- 441 **PATI:** Polar auxin transport inhibitor.
- 442 ***Physcomitrella*:** A model moss.
- 443 **PIN:** An auxin efflux facilitator class.
- 444 **Plasmodesmata:** Plasma membrane lined channels that can provide cytoplasmic continuity between plant
445 cells.
- 446 **Primordium:** An organ or tissue at the earliest stages of development.
- 447 ***Selaginella*:** A model lycophyte.
- 448 **Sporangium:** A plant organ in which meiosis occurs to make spores.
- 449 **Sporophyte:** A stage of the land plant life cycle during which meiosis occurs.
- 450
- 451
- 452
- 453
- 454

